



## Classification of chronic kidney disease using GFR and ACR categories

GFR and ACR categories and risk of adverse outcomes			ACR categories (mg/mmol), description and range		
			<3 Normal to mildly increased	3–30 Moderately increased	>30 Severely increased
			A1	A2	A3
GFR categories (ml/min/1.73 m <sup>2</sup> ), description and range	≥90 Normal and high	G1	No CKD in the absence of markers of kidney damage		
	60–89 Mild reduction related to normal range for a young adult	G2			
	45–59 Mild–moderate reduction	G3a <sup>1</sup>			
	30–44 Moderate–severe reduction	G3b			
	15–29 Severe reduction	G4			
	<15 Kidney failure	G5			


  
**Increasing risk**


  
**Increasing risk**

<sup>1</sup> Consider using eGFR<sub>cystatinC</sub> for people with CKD G3aA1 (see recommendations 1.1.14 and 1.1.15)

Abbreviations: ACR, albumin:creatinine ratio; CKD, chronic kidney disease; GFR, glomerular filtration rate

Adapted with permission from Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group (2013) KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney International (Suppl. 3): 1–150*